


**CASIO®**

CASIO  
TEACHING  
MATERIALS

# BASIC MATHEMATICS EXERCISE

with fx-991CW



**DIGEST EDITION**

Download  
the full version here

# Basic Mathematics Exercise with fx-991CW

### Introduction

These teaching materials were created with the hope of conveying to many teachers and students the appeal of scientific calculators.

#### **(1) Change awareness (emphasizing the thinking process) and boost efficiency in learning mathematics**

- By reducing the time spent on manual calculations, we can have learning with a focus on the thinking process that is more efficient.
- This reduces the aversion to mathematics caused by complicated calculations, and allows students to experience the joy of thinking, which is the essence of mathematics.

#### **(2) Diversification of learning materials and problem-solving methods**

- Making it possible to do difficult calculations manually allows for diversity in learning materials and problem-solving methods.

#### **(3) Promoting understanding of mathematical concepts**

- By using the various functions of the scientific calculator in creative ways, students are able to deepen their understanding of mathematical concepts through calculations and discussions from different perspectives than before.
- This allows for exploratory learning through easy trial and error of questions.
- Listing and graphing of numerical values by means of tables allows students to discover laws and to understand visually.

### Features of these teaching materials

- Makes classes more efficient by using scientific calculators
- Covers most mathematics topics in lower and upper secondary schools (17 units, 294 sections)
- Enables even beginners to confidently use scientific calculators in class
- Worksheets are also provided for student practice
- Helps students to make full use of scientific calculator's functions in each unit
- Operational instructions with key log display



**Better Mathematics Learning  
with Scientific Calculator**

# Linear inequalities (2)

## TARGET

To understand how to solve linear inequalities.

## STUDY GUIDE

### How to solve linear inequalities

Mathematical theorems and concepts are explained in detail.  
A scientific calculator is used to check and derive formulas according to the topic.

An inequality expressing a range of possible values for  $x$  is called an **inequality** for  $x$ . The range of values of  $x$  that satisfy the inequality for  $x$  is called the **solution of the inequality**. Finding all the solutions to an inequality is called **solving the inequality**. When all the terms of an inequality are arranged on the left side, such as "(linear expression of  $x$ )  $\geq 0$ " or "(linear expression of  $x$ )  $< 0$ ", then the left side is an inequality expressed as a linear expression, which is called a **linear inequality**.

**Ex.** How to solve the linear inequality  $3x - 5 < 7$

From the properties of inequalities that if  $a < b$  then  $a + c < b + c$ , we get  $3x - 5 + 5 < 7 + 5$  and  $3x < 12$  ... (i)

From the properties of inequalities that when  $m > 0$ , if  $a < b$  then  $\frac{a}{m} < \frac{b}{m}$ , we get  $\frac{3x}{3} < \frac{12}{3}$  and  $x < 4$  ... (ii)

So, in the expression derived in (i) by the above transformation, we could consider moving  $-5$ , from  $3x - 5 < 7$ , to the right side, which changed the sign to  $+5$ , so we could derive  $3x < 7 + 5$ . In other words, the inequality can be solved by transposition, **the same as when solving equations**.

## EXERCISE

Students learn basic examples based on the explanation in STUDY GUIDE.



1 Solve the following linear inequalities.

(1)  $4x - 3 < 2x + 9$

Transpose to get  $4x - 2x < 9 + 3$

Arrange to get  $2x < 12$

Divide both sides by 2 to get  $x < 6$

check

Explains how to use the scientific calculator to solve problems and check answers.

$$x < 6$$

On the scientific calculator, use the Table function to compare both sides of the inequality.

Press  $\odot$ , select [Table], press  $\odot$ , then clear the previous data by pressing  $\odot$

Press  $\odot$ , select [Define  $f(x)/g(x)$ ], press  $\odot$ ,

select [Define  $f(x)$ ], press  $\odot$

After inputting  $f(x) = 4x - 3$ , press  $\odot$

In the same way, input  $g(x) = 2x + 9$ .

Press  $\odot$ , select [Table Range], press  $\odot$


After inputting [Start:4, End:7, Step:1], select [Execute], press  $\odot$

Check the start and end of the solution.

When  $x < 6$ , then  $4x - 3 < 2x + 9$ ,

when  $x = 6$ , then  $4x - 3 = 2x + 9$ ,

when  $x > 6$ , then  $4x - 3 > 2x + 9$ , so each can be checked.



$$f(x) = 4x - 3$$

$$g(x) = 2x + 9$$

Table Range			
Start:	4		
End:	7		
Step:	1		

	$f(x)$	$g(x)$
1	1	12
2	5	17
3	9	21
4	13	25

The content of this page is part of "1. Algebraic Expressions and Linear Functions".

(2)  $5x+6 \geq 8x-9$

Transpose to get  $5x-8x \geq -9-6$

Arrange to get  $-3x \geq -15$

Divide both sides by  $-3$  to get  $x \leq 5$

$x \leq 5$

check

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

Press  $\text{2ND}$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ ,

select [Define  $f(x)$ ], press  $\text{OK}$

After inputting  $f(x)=5x+6$ , press  $\text{EXE}$

In the same way, input  $g(x)=8x-9$ .

Press  $\text{2ND}$ , select [Table Range], press  $\text{OK}$

After inputting [Start:3, End:6, Step:1], select [Execute], press  $\text{EXE}$

When  $x < 5$ , then  $5x+6 > 8x-9$ ,

when  $x = 5$ , then  $5x+6 = 8x-9$ ,

when  $x > 5$ , then  $5x+6 < 8x-9$ , so each can be checked.

$f(x)=5x+6$

$g(x)=8x-9$

Table Range  
Start:3  
End:6  
Step:1

x	f(x)	g(x)
1	11	-1
2	16	7
3	21	15
4	26	23
5	31	31
6	36	39

(3)  $\frac{1}{3}x - 1 \leq \frac{1}{2}x + 1$

Multiply both sides by 6 to get  $\left(\frac{1}{3}x - 1\right) \leq \left(\frac{1}{2}x + 1\right)$

Remove the brackets to get  $2x-6 \leq 3x+6$

Transpose to get  $2x-3x \leq 6+6$

Arrange to get  $-x \leq 12$

Divide both sides by  $-1$  to get  $x \geq -12$

[Table] is one of the distinctive functions of the fx-991CW that is useful for summarizing a large number of values in a table for easy viewing, or for finding some kind of rule from the created table. The fx-991CW allows students to input two functions,  $f(x)$  and  $g(x)$ . Here, they can use [Table] to examine how the values on both sides of a linear inequality change. This process is also useful for understanding the nature of linear inequalities.

check

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

Press  $\text{2ND}$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$

After inputting  $f(x)=\frac{1}{3}x - 1$ , press  $\text{EXE}$

In the same way, input  $g(x)=\frac{1}{2}x + 1$ .

Press  $\text{2ND}$ , select [Table Range], press  $\text{OK}$

After inputting [Start:-13, End:-10, Step:1], select [Execute], press  $\text{EXE}$

When  $x < -12$ , then  $\frac{1}{3}x - 1 > \frac{1}{2}x + 1$ ,

when  $x = -12$ , then  $\frac{1}{3}x - 1 = \frac{1}{2}x + 1$ ,

when  $x > -12$ , then  $\frac{1}{3}x - 1 < \frac{1}{2}x + 1$ , so each can be checked.

$f(x)=\frac{1}{3}x-1$

$g(x)=\frac{1}{2}x+1$

Table Range  
Start:-13  
End:-10  
Step:1

x	f(x)	g(x)
1	-13.333	-5.5
2	-12.666	-5
3	-12	-4.5
4	-11.333	-4

-13

Where to use the scientific calculator.



- 2] If 1 piece of item A costs \$8 and 1 piece of item B costs \$5, and you want to buy 10 pieces at a total price of less than or equal to \$62, then what is the maximum number of pieces of item A that you should buy?

Let the number of pieces of item A be  $x$  pieces.

The number of pieces of item B is  $(10-x)$  pieces, so the total cost is  $\{8x+5(10-x)\}$ .

Since it is less than or equal to \$62, we get  $8x+5(10-x) \leq 62$ .

So, the answer is  $8x+50-5x \leq 62$ ,  $8x-5x \leq 62-50$ ,  $3x \leq 12$ , so  $x \leq 4$

Therefore, you should buy 4 or fewer pieces of item A.

### 4 or fewer pieces

#### check

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

Press  $\text{MODE}$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ ,

select [Define  $f(x)$ ], press  $\text{OK}$

After inputting  $f(x)=8x+5(10-x)$ , press  $\text{EXE}$

In the same way, input  $g(x)=62$ .

Press  $\text{MODE}$ , select [Table Range], press  $\text{OK}$

After inputting [Start:0, End:4, Step:1], select [Execute], press  $\text{EXE}$

When  $x \leq 4$ , then we can confirm that  $8x+5(10-x) \leq 62$ .

$$f(x)=8x+5(10-x)$$

$$g(x)=62$$

Table Range  
Start:0  
End :4  
Step :1

$x$	$f(x)$	$g(x)$
1	53	62
2	56	62
3	59	62
4	62	62

## PRACTICE

Students can do practice problems similar to those in EXERCISE. They can also practice using the scientific calculator as they learned to in check. In PRACTICE and ADVANCED the answers are printed in red. (Separate data is also available without the red parts, so it can be used for exercises.)



◆ Solve the following linear inequ

(1)  $7x-4 < 4x+8$

**Transpose to get  $7x-4x < 8+4$**

**Arrange to get  $3x < 12$**

**Divide both sides by 3 to get  $x < 4$**

$x < 4$

check

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

Press  $\text{MODE}$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$

After inputting  $f(x)=7x-4$ , press  $\text{EXE}$

In the same way, input  $g(x)=4x+8$ .

Press  $\text{MODE}$ , select [Table Range], press  $\text{OK}$

After inputting [Start:2, End:5, Step:1], select [Execute], press  $\text{EXE}$

When  $x < 4$ , then  $7x-4 < 4x+8$ ,

when  $x=4$ , then  $7x-4=4x+8$ ,

when  $x > 4$ , then  $7x-4 > 4x+8$ , so each can be checked.

f(x)=7x-4	
1	3
2	10
3	17
4	24
5	31

g(x)=4x+8	
1	12
2	16
3	20
4	24
5	28

Table Range	
Start:	2
End:	5
Step:	1

x	f(x)	g(x)
1	3	12
2	10	16
3	17	20
4	24	24
5	31	28

2

(2)  $2x+5 \geq 3x-2$

**Transpose to get  $2x-3x \geq -2-5$**

**Arrange to get  $-x \geq -7$**

**Divide both sides by  $-1$  to get  $x \leq 7$**

$x \leq 7$

check

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

Press  $\text{MODE}$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$

After inputting  $f(x)=2x+5$ , press  $\text{EXE}$

In the same way, input  $g(x)=3x-2$ .

Press  $\text{MODE}$ , select [Table Range], press  $\text{OK}$

After inputting [Start:6, End:9, Step:1], select [Execute], press  $\text{EXE}$

When  $x < 7$ , then  $2x+5 > 3x-2$ ,

when  $x=7$ , then  $2x+5=3x-2$ ,

when  $x > 7$ , then  $2x+5 < 3x-2$ , so each can be checked.

f(x)=2x+5	
1	7
2	9
3	11
4	13
5	15

g(x)=3x-2	
1	1
2	4
3	7
4	10
5	13

Table Range	
Start:	6
End:	9
Step:	1

x	f(x)	g(x)
1	7	1
2	9	4
3	11	7
4	13	10
5	15	13
6	17	16
7	19	19
8	21	22
9	23	25

6

## PRACTICE

- 1 Determine the axes and vertexes of the graph of the quadratic function  $y = x^2 - 8x + 10$ .

$$y = x^2 - 8x + 10 = (x - 4)^2 - 4^2 + 10 = (x - 4)^2 - 6$$

Therefore, we get an axis of  $x=4$  and a vertex of  $(4, -6)$ .



$$x=4, (4, -6)$$

check




Press  $\odot$ , select [Equation], press  $\text{OK}$

Select [Polynomial], press  $\text{OK}$ , select  $[ax^2 + bx + c]$ , press  $\text{OK}$

$\text{1}$   $\text{EXE}$   $\text{—}$   $\text{8}$   $\text{EXE}$   $\text{1}$   $\text{0}$   $\text{EXE}$   $\text{EXE}$



$\text{EXE}$   $\text{EXE}$   $\text{EXE}$  Press  $\uparrow$   $\odot$ , scan the QR code to display a graph.



Students can generate a QR code based on the calculation results and data displayed on the scientific calculator, and then scan it with a smartphone or tablet to display a graph. This is one of the distinctive features of the fx-991CW. Here, a graph of  $y = x^2 - 8x + 10$  is drawn to help students understand visually that the vertex of the graph is  $(4, -6)$ .



The content of this page is part of "2. Quadratic Functions".



2 Draw the graphs of the following functions.

(1)  $y = 2x^2 - 8x + 7$

$$y = 2x^2 - 8x + 7 = 2(x^2 - 4x) + 7$$

$$= 2\{(x - 2)^2 - 2^2\} + 7 = 2(x - 2)^2 - 1$$


Therefore, we get an axis of  $x=2$  and a vertex of  $(2, -1)$ .

(2)  $y = -3x^2 + 9x - 5$

$$y = -3x^2 + 9x - 5 = -3(x^2 - 3x) - 5$$

$$= -3\left\{x - \frac{3}{2}\right\}^2 - \left(\frac{3}{2}\right)^2 - 5 = -3\left(x - \frac{3}{2}\right)^2 + \frac{7}{4}$$

Therefore, we get an axis of  $x = \frac{3}{2}$  and a vertex of  $\left(\frac{3}{2}, \frac{7}{4}\right)$ .



check

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

Press  $\text{MODE}$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$ , after inputting  $f(x) = 2x^2 - 8x + 7$ , press  $\text{EXE}$



$f(x) = 2x^2 - 8x + 7$

In the same way, input  $g(x) = -3x^2 + 9x - 5$ .


Press  $\text{MODE}$ , select [Table Range], press  $\text{OK}$ , after inputting [Start:-5, End:5, Step:1], select [Execute], press  $\text{EXE}$

$g(x) = -3x^2 + 9x - 5$

Table Range  
Start:-5  
End:5  
Step:1

Press  $\uparrow$   $\text{X}$ , scan the QR code to display a graph.

	$x$	$f(x)$	$g(x)$
1	-5	97	-125
2	-4	71	-89
3	-3	49	-59
4	-2	31	-35



By using [Table], students can show the points that a graph passes through on a table. By plotting these points, they can even easily draw the graph freehand.

In addition, students can use a QR code to confirm whether the graph they have drawn is correct.





- 3 We know that when an object is thrown vertically upward from a height of 0 m at a velocity of  $v$  m/s, the relation between the height  $y$  m of the object and the time  $x$  seconds since it was thrown is  $y = -4.9x^2 + vx$ . So, when a ball is thrown vertically from a height of 0 m at a velocity of 9.8 m/s, find the time it takes to reach the highest point and the height of that highest point. Note that the air resistance can be ignored.

$$y = -4.9x^2 + 9.8x = -4.9(x^2 - 2x) = -4.9\{(x - 1)^2 - 1^2\} = -4.9(x - 1)^2 + 4.9$$

Therefore, the graph of this function is convex upward, the axis is  $x=1$ , and the vertex is (1, 4.9).

Thus, the ball takes 1 second to reach its highest point, and the height of that highest point is 4.9 m.


1 second, 4.9 m

check




Press  $\odot$ , select [Equation], press  $\odot$

Select [Polynomial], press  $\odot$ , select [ $ax^2 + bx + c$ ], press  $\odot$

$\ominus$  4  $\odot$  9  $\odot$  9  $\odot$  8  $\odot$  0  $\odot$   $\odot$



$\odot$   $\odot$   $\odot$  Press  $\uparrow$   $\odot$ , scan the QR code to display a graph.



Real-life problems are also described. In this question, students can use what they learned in PRACTICE two pages earlier.

At the vertex of the graph,  $y$  (height) starts to decrease, so we know the instant that the ball begins to fall. Note that the graph is not the trajectory of the ball.

# Finding an angle

## TARGET

To understand how to use 2 sides of a right-angled triangle to find the angles of the right-angled triangle.

Students can identify the objective to learn in each section.

## STUDY GUIDE

### Inverse functions of trigonometric ratio (inverse trigonometric functions)

Angles can be found by using the values of the trigonometric ratio.



**Ex.** When  $\sin A = \frac{1}{3}$ , angle  $A$  is expressed as follows.


$A = \sin^{-1} \frac{1}{3}$  Simple examples on how to apply equations and theorems.

$$\sin A = \frac{1}{3} \Rightarrow A = \sin^{-1} \frac{1}{3} = 19.47^\circ$$

$$\left( \text{*Namely } \sin 19.47^\circ = \frac{1}{3} \right)$$

$$\text{In the same way, from } \cos A = \frac{2\sqrt{2}}{3}, \tan A = \frac{1}{2\sqrt{2}}$$

$$\text{we can show } A = \cos^{-1} \frac{2\sqrt{2}}{3}, A = \tan^{-1} \frac{1}{2\sqrt{2}}.$$



Calculator display:  $\sin^{-1}\left(\frac{1}{3}\right)$   
19.47122063


**19.47°**

## EXERCISE



Find the measure of  $A$  to the 2nd decimal place for the  $\triangle ABC$  below.

(1)



$$\text{When } \sin A = \frac{3}{5}, \text{ we get } A = \sin^{-1} \frac{3}{5} = 36.87^\circ.$$

Calculator display:  $\sin^{-1}\left(\frac{3}{5}\right)$   
36.86989765

**36.87°**

**OTHER METHODS** From  $\cos A = \frac{4}{5}$ , we get  $A = \cos^{-1} \frac{4}{5} = 36.87^\circ$ .

Calculator display:  $\cos^{-1}\left(\frac{4}{5}\right)$   
36.86989765


**OTHER METHODS** From  $\tan A = \frac{3}{4}$ , we get  $A = \tan^{-1} \frac{3}{4} = 36.87^\circ$ .

Calculator display:  $\tan^{-1}\left(\frac{3}{4}\right)$   
36.86989765

Alternative solutions and different verification methods for previously presented problems.

The content of this page is part of "3. Trigonometry".

(2)



From  $\sin A = \frac{1}{2}$ , we get  $A = \sin^{-1} \frac{1}{2} = 30^\circ$ .

↑ sin 1 = 2 > ) EXE

$$\sin^{-1}\left(\frac{1}{2}\right)$$

30

**30°**

**OTHER METHODS**

From  $\cos A = \frac{\sqrt{3}}{2}$ , we get  $A = \cos^{-1} \frac{\sqrt{3}}{2} = 30^\circ$ .

↑ cos √ 3 2 > ) EXE

$$\cos^{-1}\left(\frac{\sqrt{3}}{2}\right)$$

30

**OTHER METHODS**

From  $\tan A = \frac{1}{\sqrt{3}}$ , we get  $A = \tan^{-1} \frac{1}{\sqrt{3}} = 30^\circ$ .

↑ tan 1 √ 3 > > ) EXE

$$\tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$$


30

## PRACTICE



Find the measure of  $A$  to the 2nd decimal place for the  $\triangle ABC$  below.

(1)



From  $\sin A = \frac{5}{13}$ , we get  $A = \sin^{-1} \frac{5}{13} = 22.62^\circ$ .

↑ sin 5 = 13 > ) EXE


$$\sin^{-1}\left(\frac{5}{13}\right)$$

22.61986495

**22.62°**

Key operations are carefully explained to students who are using a scientific calculator for the first time.

(2)



From  $\sin A = \frac{7}{11}$ , we get  $A = \sin^{-1} \frac{7}{11} = 39.52^\circ$ .

↑ sin 7 = 11 > ) EXE

$$\sin^{-1}\left(\frac{7}{11}\right)$$

39.52119636

**39.52°**

Tutorial videos are also available for students who are new to the fx-991CW. Be sure to view these!



# Trigonometric ratio formula including tangent

## TARGET


To understand a trigonometric ratio formula including tangent.

## STUDY GUIDE

### Correlation of trigonometric ratios (2)

The following relations are derived from the right-angled triangle shown in the diagram on the right.

$$(1) \frac{\sin A}{\cos A} = \tan A \quad (2) 1 + \tan^2 A = \frac{1}{\cos^2 A}$$



#### explanation

- (1) From the right-angled triangle shown in the diagram on the right

$$\frac{\sin A}{\cos A} = \frac{\frac{a}{c}}{\frac{b}{c}} = \frac{a}{b} = \tan A$$

$$\Rightarrow \frac{\sin A}{\cos A} = \tan A$$

#### EXTRA Info.


Calculating fractions

$$\frac{\frac{A}{B}}{\frac{C}{D}} = \frac{A \times D}{B \times C}$$

Proof  $\frac{\frac{A}{B}}{\frac{C}{D}} = \frac{A}{B} \div \frac{C}{D} = \frac{A}{B} \times \frac{D}{C} = \frac{A \times D}{B \times C}$

Ex.  $\frac{\frac{3}{5}}{\frac{7}{4}} = \frac{3 \times 4}{5 \times 7} = \frac{12}{35}$

Press  $\odot$ , select [Calculate], press  $\text{OK}$



- (2) Divide both sides of  $\sin^2 A + \cos^2 A = 1$  by  $\cos^2 A$

such that  $\frac{\sin^2 A}{\cos^2 A} + \frac{\cos^2 A}{\cos^2 A} = \frac{1}{\cos^2 A}$

$$\left( \frac{\sin A}{\cos A} \right)^2 + 1 = \frac{1}{\cos^2 A}$$

$$\Rightarrow \tan^2 A + 1 = \frac{1}{\cos^2 A}$$

The content of this page is part of "3. Trigonometry".



## Proofs and checks of mathematical formulas.

proof

(1) Calculate the value of the function to confirm the formula (1).

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\odot$

Press  $\odot\odot$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$

Input  $f(x) = \frac{\sin x}{\cos x}$ .

$\sin(x) \div \cos(x) \text{ EXE}$

$$f(x) = \frac{\sin(x)}{\cos(x)}$$

Press  $\odot\odot$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $g(x)$ ], press  $\text{OK}$

Input  $g(x) = \tan x$ .

$\tan(x) \text{ EXE}$

$$g(x) = \tan(x)$$

Press  $\odot\odot$ , select [Table Range], press  $\text{OK}$ , after inputting [Start:0, End:90, Step:5], select [Execute], press  $\text{EXE}$

Table Range  
End :90  
Step :5  
Execute

We can confirm that  $\frac{\sin A}{\cos A} = \tan A$  holds regardless of the value of  $A$ .

x	f(x)	g(x)
0	0	0
5	0.0874	0.0874
10	0.1763	0.1763
15	0.2679	0.2679

When  $x=0$ , then  $f(x)=g(x)=0$

When  $x=5$ , then  $f(x)=g(x)=0.0874$

As shown below, for any  $x$ , we get  $f(x)=g(x)$ .

(2) Calculate the value of the function to confirm the formula (2).

Press  $\odot$  to clear the previous data, press  $\odot\odot$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$

Input  $f(x) = 1 + \tan^2 x$ .

$1 + \tan(x)^2 \text{ EXE}$

$$f(x) = 1 + \tan(x)^2$$

Press  $\odot\odot$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $g(x)$ ], press  $\text{OK}$

Input  $g(x) = \frac{1}{\cos^2 x}$ .

$1 \div \cos(x)^2 \text{ EXE}$

$$g(x) = \frac{1}{\cos(x)^2}$$

Press  $\odot\odot$ , select [Table Range], press  $\text{OK}$ , after inputting [Start:0, End:90, Step:5], select [Execute], press  $\text{EXE}$

Table Range  
End :90  
Step :5  
Execute

We can confirm that  $1 + \tan^2 A = \frac{1}{\cos^2 A}$  holds regardless of the value of  $A$ .

x	f(x)	g(x)
0	1	1
5	1.0076	1.0076
10	1.031	1.031
15	1.0717	1.0717

On this page, students are using [Table] to check what the formula is asserting. Specifically, for each of the formulas (1) and (2) on the previous page, the students are checking that  $f(x) = g(x)$  holds for any value of  $x$ , with  $f(x)$  on the left side and  $g(x)$  on the right side. This is an effective way to correctly understand the meaning of a new formula.

# Uses of exponential functions

## TARGET

To understand how to solve problems using graphs of exponential functions.

## EXERCISE

Real-life problems are also described.



Suppose we examined the transmission of an infectious disease every day, and our results showed that the number of newly infected people increased by  $r$  times after 1 day. For example, on a given day 100 people are infected, and if  $r=1.5$ , then after 1 day the number of newly infected people would be  $100 \times 1.5 = 150$  (people), and after 2 days it would be  $150 \times 1.5 = 225$  (people). Calculating this from 2 days ago, gives us  $100 \times 1.5^2$  (people). From this, we can realize the following. That after  $x$  days, the number of newly infected people will be  $y$  times the initial number of people, which we can express as  $y = r^x$ . Now, solve the following problems.

- (1) Given that  $r=1.5$  does not change. Find how many times the number of people infected on the first day would be newly infected after 10 days. Also, given that 100 people were infected on the first day, find how many people would be newly infected after 10 days.

Substitute  $x=10$  into  $y = 1.5^x$ , such that  $y = 1.5^{10} = 57.66 \dots$  (times).

The number of people newly infected after 10 days would be

$100 \times 57.66 \dots = 5766 \dots$  (people).

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

Press  $\text{MODE}$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$


After inputting  $f(x) = 1.5^x$ , press  $\text{EXE}$

Press  $\text{MODE}$ , select [Table Range], press  $\text{OK}$

After inputting [Start:0, End:10, Step:1], select [Execute], press  $\text{EXE}$

$x$	$f(x)$	$g(x)$
7	17.085	ERROR
8	25.628	ERROR
9	38.443	ERROR
10	57.663	ERROR

57.66503906



**Approximately 58 times, approximately 5800 people**

- (2) Find a value for  $r$  that would reduce the number of newly infected people.

For example, examine what happens when  $r=0.8$  and  $0.4$ .

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

Press  $\text{MODE}$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$ , after inputting  $f(x) = 0.8^x$ , press  $\text{EXE}$


In the same way, input  $g(x) = 0.4^x$ .

Press  $\text{MODE}$ , select [Table Range], press  $\text{OK}$

After inputting [Start:0, End:10, Step:1], select [Execute], press  $\text{EXE}$

$x$	$f(x)$	$g(x)$
7	0.2097	$1.6 \times 10^{-2}$
8	0.1677	$6.5 \times 10^{-4}$
9	0.1342	$2.6 \times 10^{-4}$
10	0.1074	$1 \times 10^{-4}$

0.1073741824



When  $0 < r < 1$ , the curve falls to the right, so we know it is decreasing.

$0 < r < 1$

The content of this page is part of "5. Exponential and Logarithmic Functions".



# Distance between points and lines

## TARGET

To understand about distances between points and lines.

## STUDY GUIDE

### Distance between points and lines



When we let  $H$  be the intersection of a perpendicular line drawn from point  $A$  to the line  $l$ , then the length  $d$  of  $AH$  is called the **distance** between point  $A$  and the line  $l$ , which we can find as follows.

- (1)  $d$  is the distance between  $A(x_1, y_1)$  and line  $ax+by+c=0$

$$d = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$$

- (2)  $d$  is the distance between the origin  $O$  and line  $ax+by+c=0$

$$d = \frac{|c|}{\sqrt{a^2 + b^2}}$$



Formulas and their supplementary explanations.

#### explanation

Let the points be  $A(x_1, y_1)$  and  $H(x_2, y_2)$ , such that  $d = AH = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$  ... (i)

Since the line  $AH$  is perpendicular to the line  $ax+by+c=0$ , we get  $bx - ay + c' = 0$  ... (ii).

It passes through point  $A(x_1, y_1)$ , so we get  $bx_1 - ay_1 + c' = 0$ ,  $c' = -bx_1 + ay_1$

By substituting this into (ii), we get  $bx - ay - bx_1 + ay_1 = 0$ , so  $b(x - x_1) - a(y - y_1) = 0$

The line  $AH$  passes through  $H(x_2, y_2)$ , so we get  $b(x_2 - x_1) - a(y_2 - y_1) = 0$  ... (iii)

Since point  $H(x_2, y_2)$  is a point on the line  $ax+by+c=0$ , we get  $ax_2 + by_2 + c = 0$

By transforming this, we get  $a(x_2 - x_1) + b(y_2 - y_1) + ax_1 + by_1 + c = 0$  ... (iv)

If we solve (iii) and (iv) for  $x_2 - x_1$  and  $y_2 - y_1$ , then we get

$$x_2 - x_1 = -\frac{a(ax_1 + by_1 + c)}{a^2 + b^2}, y_2 - y_1 = -\frac{b(ax_1 + by_1 + c)}{a^2 + b^2}$$

These are substituted into (i), for

$$d = \sqrt{\left\{-\frac{a(ax_1 + by_1 + c)}{a^2 + b^2}\right\}^2 + \left\{-\frac{b(ax_1 + by_1 + c)}{a^2 + b^2}\right\}^2} = \sqrt{\frac{(a^2 + b^2)(ax_1 + by_1 + c)^2}{(a^2 + b^2)^2}} = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$$

If the point  $A$  is the origin, then  $x_1 = 0$  and  $y_1 = 0$ , so  $d = \frac{|c|}{\sqrt{a^2 + b^2}}$

The content of this page is part of  
"6. Equations of Lines and Circles".



## EXERCISE



Find the distances between the following points and lines.

- (1) Point (2, 3) and line  $5x - 4y + 8 = 0$

$$d = \frac{|5 \cdot 2 - 4 \cdot 3 + 8|}{\sqrt{5^2 + (-4)^2}} = \frac{6}{\sqrt{41}} = \frac{6\sqrt{41}}{41}$$

$$\frac{6\sqrt{41}}{41}$$

- (2) The origin and line  $2x + 3y - 9 = 0$

$$d = \frac{|-9|}{\sqrt{2^2 + 3^2}} = \frac{9}{\sqrt{13}} = \frac{9\sqrt{13}}{13}$$

$$\frac{9\sqrt{13}}{13}$$

- (3) Point (1, -5) and line  $y = 2x + 3$

By transforming  $y = 2x + 3$ , we get  $2x - y + 3 = 0$

$$d = \frac{|2 \cdot 1 - 1 \cdot (-5) + 3|}{\sqrt{2^2 + (-1)^2}} = \frac{10}{\sqrt{5}} = 2\sqrt{5}$$

$$2\sqrt{5}$$

Repeating similar calculations is one of the things that a scientific calculator is good at. By using [VARIABLE], students can see how values change when they change multiple variables (in this question, when the values of A, B, C, x, and y are changed). By trying out various values, the students are likely to notice some unexpected regularities!

## check

On the scientific calculator, use the VARIABLE function to calculate the distance between a point and a line.


Press  $\odot$ , select [Calculate], press  $\text{OK}$

Input the formula for the distance between a point and a line.




$$\frac{|Ax+By+C|}{\sqrt{A^2+B^2}}$$


- (1) In the VARIABLE screen, input [A=5, B=-4, C=8, x=2, and y=3], and then calculate.



- (2) In the same way, input [A=2, B=3, C=-9, x=0, and y=0], and then calculate.



- (3) In the same way, input [A=2, B=-1, C=3, x=1, and y=-5], and then calculate.



Use the scientific calculator to confirm that equations and inequalities hold.

## EXERCISE




Use the scientific calculator to confirm that the following equations and inequalities hold.

(1)  $a^3 + b^3 = (a + b)^3 - 3ab(a + b)$

$$a^3 + b^3 = (a + b)^3 - 3ab(a + b) \Leftrightarrow (a + b)^3 - 3ab(a + b) - (a^3 + b^3) = 0$$


Therefore, we simply need to confirm that the right side – the left side = 0 (or, the left side – the right side = 0).

Press  $\odot$ , select [Calculate], press  $\text{OK}$



Input  $(A + B)^3 - 3AB(A + B) - (A^3 + B^3)$ .



( ( 4 + 5 )  $\square^3$  3 > - 3 ( 4 + 5 ) ( ( 4 + 5 )  $\square^3$  3 > )



Use the VARIABLE function to assign any values to A and B.

(Example) To input A=13 and B=-28

$\odot$  1 3  $\text{EXE}$  > - 2 8  $\text{EXE}$   $\rightarrow$   $\text{EXE}$



We can confirm  $(A + B)^3 - 3AB(A + B) - (A^3 + B^3) = 0$  using any values, so we can verify the validity of  $a^3 + b^3 = (a + b)^3 - 3ab(a + b)$  holds. You may want to verify other examples.


(2)  $(ac + bd)^2 \leq (a^2 + b^2)(c^2 + d^2)$

$$(ac + bd)^2 \leq (a^2 + b^2)(c^2 + d^2) \Leftrightarrow (ac + bd)^2 - (a^2 + b^2)(c^2 + d^2) \leq 0$$

Therefore, we simply need to confirm that the left side – the right side  $\leq 0$  (or, the right side – the left side  $\geq 0$ ).

Input  $(AC + BD)^2 - (A^2 + B^2)(C^2 + D^2)$ .



( ( 4 + 6 )  $\square^2$  - ( 5 + 1 )  $\square^2$  ) ( ( 4 + 5 )  $\square^2$  ) ( ( 6 + 1 )  $\square^2$  )



Use the VARIABLE function to assign any values to A, B, C, D.

(Example) To input A=5, B=-3, C=-4, and D=9

$\odot$  5  $\text{EXE}$  > - 3  $\text{EXE}$  > - 4  $\text{EXE}$  > 9  $\text{EXE}$   $\rightarrow$   $\text{EXE}$



We can confirm  $(AC + BD)^2 - (A^2 + B^2)(C^2 + D^2) < 0$  using any values, so we can verify the validity of  $(ac + bd)^2 \leq (a^2 + b^2)(c^2 + d^2)$  holds. You may want to verify other examples.

On this page, students use [VARIABLE] to verify that the equality and inequality of  $a^3 + b^3 = (a + b)^3 - 3ab(a + b)$  and  $(ac + bd)^2 \leq (a^2 + b^2)(c^2 + d^2)$  hold.

The content of this page is part of "7. Formulas and Proofs".

# Sequences

## TARGET

To understand how to express a sequence and its general term.

## STUDY GUIDE

### General terms of sequences

Numbers arranged in a row according to a rule are called a **sequence**, and each individual number is called a **term**.

A sequence with a finite number of terms is called a **finite sequence**, and a sequence with an infinite number of terms is called an **infinite sequence**.

We say that a finite sequence has a **number of terms** and there is a **first term** and a **last term**.

Sequences are generally expressed as follows using letters with indices.

$$a_1, a_2, a_3, \dots, a_n, \dots$$

The sequence above can also be expressed as  $\{a_n\}$ .

When the  $n$ th term in the sequence  $\{a_n\}$  is expressed in an expression as  $n$ , we say it is the **general term** of the sequence  $\{a_n\}$ .

**Ex.** Given the sequence  $\{a_n\}$  of 2, 4, 6, 8, and 10 has 5 terms:  $a_1=2$ (first term),  $a_2=4$ ,  $a_3=6$ ,  $a_4=8$ ,  $a_5=10$ (last term)

From  $a_1=2 \cdot 1$ ,  $a_2=2 \cdot 2$ ,  $a_3=2 \cdot 3$ ,  $a_4=2 \cdot 4$ , and  $a_5=2 \cdot 5$ , we can estimate that the general term is  $a_n=2n$ .

## EXERCISE



- 1 Find the first to 4th terms of the sequence  $\{a_n\}$  whose general term is expressed by the following expression.

(1)  $a_n = 3n + 2$

When the first term is  $n=1$ , we get  $a_1=3 \cdot 1 + 2=5$

When the 2nd term is  $n=2$ , we get  $a_2=3 \cdot 2 + 2=8$

When the 3rd term  $n=3$ , we get  $a_3=3 \cdot 3 + 2=11$

When the 4th term is  $n=4$ , we get  $a_4=3 \cdot 4 + 2=14$

$$\underline{a_1 = 5, a_2 = 8, a_3 = 11, a_4 = 14}$$

### check

On the scientific calculator, use the Table function to confirm each term of the sequence.

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\odot$




Press  $\odot$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$

After inputting  $f(x)=3x+2$ , press  $\text{EXE}$

Press  $\odot$ , select [Table Range], press  $\text{OK}$

After inputting [Start:1, End:4, Step:1], select [Execute],

press  $\text{EXE}$



The content of this page is part of  
"10. Sequences".



# Recurrence formula (1)

## EXERCISE



- 1 Find the 2nd to 5th terms given the first term and recurrence formula are as follows.

(1)  $a_1=2, a_{n+1}=4a_n+5$

$$a_2=4a_1+5=4\cdot 2+5=13, a_3=4a_2+5=4\cdot 13+5=57, a_4=4a_3+5=4\cdot 57+5=233, a_5=4a_4+5=4\cdot 233+5=937$$

$$\underline{a_2=13, a_3=57, a_4=233, a_5=937}$$

(2)  $a_1=5, a_{n+1}=2a_n-3$

$$a_2=2a_1-3=2\cdot 5-3=7, a_3=2a_2-3=2\cdot 7-3=11, a_4=2a_3-3=2\cdot 11-3=19, a_5=2a_4-3=2\cdot 19-3=35$$

$$\underline{a_2=7, a_3=11, a_4=19, a_5=35}$$

(3)  $a_1=1, a_{n+1}=a_n^2+1$

$$a_2=a_1^2+1=1^2+1=2, a_3=a_2^2+1=2^2+1=5, a_4=a_3^2+1=5^2+1=26, a_5=a_4^2+1=26^2+1=677$$

$$\underline{a_2=2, a_3=5, a_4=26, a_5=677}$$

## check

Press  $\odot$ , select [Spreadsheet], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

- (1) After inputting [A1:2], press  $\text{EXE}$

Press  $\text{F9}$ , select [Fill Formula], press  $\text{OK}$

After inputting [Form= $4A1+5$ ], press  $\text{EXE}$

After inputting [Range:A2:A5], press  $\text{EXE}$ , select [Confirm], press  $\text{OK}$

A	B	C	D
2			

Fill Formula  
Form =4A1+5  
Range :A2:A5  
Confirm

A	B	C	D
2			
13			
57			
233			

=4A1+5

- (2) When the sheet is displayed, move to [B1].

After inputting [B1:5], press  $\text{EXE}$

Press  $\text{F9}$ , select [Fill Formula], press  $\text{OK}$

After inputting [Form= $2B1-3$ ], press  $\text{EXE}$

After inputting [Range:B2:B5], press  $\text{EXE}$ , select [Confirm], press  $\text{OK}$

A	B	C	D
2			
13			
57			
233			

Fill Formula  
Form =2B1-3  
Range :B2:B5  
Confirm

A	B	C	D
2			
13	5		
57	7		
233	11		

=2B1-3

- (3) When the sheet is displayed, move to [C1].

After inputting [C1:1], press  $\text{EXE}$

Press  $\text{F9}$ , select [Fill Formula], press  $\text{OK}$

After inputting [Form= $C1^2+1$ ], press  $\text{EXE}$

After inputting [Range:C2:C5], press  $\text{EXE}$ , select [Confirm], press  $\text{OK}$

A	B	C	D
2			
13			
57			
233			

Fill Formula  
Form =C1^2+1  
Range :C2:C5  
Confirm

A	B	C	D
2			
13	5	1	
57	7	2	
233	11	5	

=C1^2+1

On this page, the student is introduced to using [Spreadsheet] to find the value of each term from the recurrence formula of a sequence of numbers. [Spreadsheet] makes it possible to perform calculations using a 45-row  $\times$  5-column spreadsheet.

 2 Find the general term the sequence  $\{a_n\}$  given the first term and recurrence formula are as follows.

(1)  $a_1=7, a_{n+1}=a_n-3$

The sequence  $\{a_n\}$  is an arithmetic progression with a first term of 7 and a common difference of  $-3$ ,  
so  $a_n=7+(n-1)\cdot(-3)=-3n+10$

$$\underline{a_n = -3n + 10}$$

(2)  $a_1=5, a_{n+1}=2a_n$

The sequence  $\{a_n\}$  is a geometric progression with a first term of 5 and a common ratio of 2, so  $a_n = 5 \cdot 2^{n-1}$

$$\underline{a_n = 5 \cdot 2^{n-1}}$$

(3)  $a_1=3, a_{n+1}=a_n+2n$

From  $a_{n+1}-a_n=2n$ , then for the progression of differences  $\{b_n\}$  of the sequence  $\{a_n\}$ , we get  $b_n=2n$ .

Given  $n \geq 2$ , then  $a_n = a_1 + \sum_{k=1}^{n-1} b_k = 3 + \sum_{k=1}^{n-1} 2k = 3 + 2 \cdot \frac{1}{2}(n-1)n = n^2 - n + 3$

This also holds when  $n=1$  and  $1^2 - 1 + 3 = 3$ .

Therefore, the general term is  $a_n = n^2 - n + 3$

$$\underline{a_n = n^2 - n + 3}$$

#### check

Press  $\odot$ , select [Spreadsheet], press  $\text{OK}$ , then clear the previous data by pressing  $\uparrow$

After inputting [A1:1, A2:2, A3:3, and A4:4] respectively, press  $\text{EXE}$ , move to [B1].

(1) After inputting [B1:7], press  $\text{EXE}$

Press  $\odot$ , select [Fill Formula], press  $\text{OK}$

After inputting [Form=B1-3], press  $\text{EXE}$

After inputting [Range:B2:B4], press  $\text{EXE}$ , select [Confirm], press  $\text{OK}$

	A	B	C	D
1	1	7		
2	2			
3	3			
4	4			

Fill Formula  
Form =B1-3  
Range :B2:B4  
Confirm

	A	B	C	D
1	1	7		
2	2	4		
3	3	1		
4	4	-2		

=B1-3

(2) When the sheet is displayed, move to [C1].

After inputting [C1:5], press  $\text{EXE}$

Press  $\odot$ , select [Fill Formula], press  $\text{OK}$

After inputting [Form=2C1], press  $\text{EXE}$

After inputting [Range:C2:C4], press  $\text{EXE}$ , select [Confirm], press  $\text{OK}$

	A	B	C	D
1	1	7	5	
2	2	4		
3	3	1		
4	4	-2		

Fill Formula  
Form =2C1  
Range :C2:C4  
Confirm

	A	B	C	D
1	1	7	5	
2	2	4	10	
3	3	1	20	
4	4	-2	40	

=2C1

(3) When the sheet is displayed, move to [D1].

After inputting [D1:3], press  $\text{EXE}$

Press  $\odot$ , select [Fill Formula], press  $\text{OK}$

After inputting [Form=D1+2A1], press  $\text{EXE}$

After inputting [Range:D2:D4], press  $\text{EXE}$ , select [Confirm], press  $\text{OK}$

	A	B	C	D
1	1	7	5	3
2	2	4	10	
3	3	1	20	
4	4	-2	40	

Fill Formula  
Form =D1+2A1  
Range :D2:D4  
Confirm

	A	B	C	D
1	1	7	5	3
2	2	4	10	5
3	3	1	20	9
4	4	-2	40	15

=D1+2A1

Press  $\uparrow$   $\text{X}$ , scan the QR code to display the data. (Continued on the next page.)

## PRACTICE



- ◆ The table on the right is the data for 2 variates  $x$  and  $y$ . Draw a scatter plot and find the correlation coefficient. Furthermore, determine if  $x$  and  $y$  have a correlation and whether it is positive or negative and strong or weak.

	$x$	$y$
A	3	4
B	8	7
C	12	11
D	21	14


The average values of  $x$  and  $y$  are  $\bar{x}$  and  $\bar{y}$ , the deviations are  $x - \bar{x}$  and  $y - \bar{y}$ , the product of the deviations is  $(x - \bar{x})(y - \bar{y})$ , and the squares of the deviations are  $(x - \bar{x})^2$  and  $(y - \bar{y})^2$ , which are calculated in order and summarized in the following table.

	$x$	$y$	$x - \bar{x}$	$y - \bar{y}$	$(x - \bar{x})(y - \bar{y})$	$(x - \bar{x})^2$	$(y - \bar{y})^2$
A	3	4	-8	-5	40	64	25
B	8	7	-3	-2	6	9	4
C	12	11	1	2	2	1	4
D	21	14	10	5	50	100	25
Total	44	36	0	0	98	174	58
Average	11	9	0	0	24.5	43.5	14.5

From  $s_{xy} = 24.5$ ,  $s_x = \sqrt{43.5} \simeq 6.6$ ,  $s_y = \sqrt{14.5} \simeq 3.8$ , we get

$$r = \frac{s_{xy}}{s_x s_y} = \frac{24.5}{6.6 \times 3.8} \simeq 0.98$$

The scatter plot is shown on the right.



The correlation coefficient is  $r=0.98$ , and  $x$  and  $y$  have a strong positive correlation.

check

Press  $\odot$ , select [Statistics], press  $\text{OK}$ , select [2-Variable], press  $\text{OK}$



1-Variable  
2-Variable

Input in the x column.

$\text{3}$   $\text{EXE}$   $\text{8}$   $\text{EXE}$   $\text{1}$   $\text{2}$   $\text{EXE}$   $\text{2}$   $\text{1}$   $\text{EXE}$

	x	y
1	3	0
2	8	0
3	12	0
4	21	0

Return to row 1 in the y column.

$\nabla$   $\triangleright$

	x	y
1	3	0
2	8	0
3	12	0
4	21	0

Input in the y column.

$\text{4}$   $\text{EXE}$   $\text{7}$   $\text{EXE}$   $\text{1}$   $\text{1}$   $\text{EXE}$   $\text{1}$   $\text{4}$   $\text{EXE}$

	x	y
1	3	7
2	8	11
3	12	14
4	21	

After inputting the final data, press  $\text{EXE}$ , select [2-Var Results], press  $\text{OK}$

2-Var Results  
Reg Results  
Statistics Calc

Standard deviation of  $x$ :  $\sigma_x = 6.6$

Standard deviation of  $y$ :  $\sigma_y = 3.8$

$\bar{x}$	=11
$\Sigma x$	=44
$\Sigma x^2$	=658
$\sigma^2 x$	=43.5
$\sigma x$	=6.595452979
$s^2 x$	=58

$\Sigma x$	=7.615773106
$n$	=4
$\bar{y}$	=9
$\Sigma y$	=36
$\Sigma y^2$	=382
$\sigma^2 y$	=14.5

$\sigma y$	=3.807886553
$s^2 y$	=19.33333333
$s y$	=4.396968653
$\Sigma xy$	=494
$\Sigma x^3$	=11528
$\Sigma x^2 y$	=8242

$\Sigma x^4$	=219394
$\min(x)$	=3
$\max(x)$	=21
$\min(y)$	=4
$\max(y)$	=14

Since covariance is not shown in the above table, it is found by  
(covariance) = (average of product) – (product of averages).

$$s_{xy} = \frac{\Sigma xy}{n} - \bar{x}\bar{y} = \frac{494}{4} - 11 \cdot 9 = 24.5$$


If students try to calculate the mean, variance, standard deviation, and other values by hand, it takes a very long time and they might make mistakes in their calculations. However, one of the fx-991CW's unique features is [Statistics], which students can use to find these values quickly, easily, and accurately. This allows students to focus on the process of analyzing the data and reading the trends in the data.



Select the QR code function.



Read the QR code with your smartphone.



# Events and probability

## TARGET

To understand the basics of probability.

## STUDY GUIDE

### Events and probability

#### Trials and events

An experiment or observation that can be repeated under the same conditions, such as rolling dice, is called a **trial**. The result of a trial is called an **event**, which is often expressed using a set. Also, an event that always happens in a trial is called a **sure event** for that trial, and it is expressed by  $U$ . Furthermore, an event that cannot be further divided is called a **root event**.

**Ex.** The sure events in a trial rolling 1 die is  $U = \{1, 2, 3, 4, 5, \text{ and } 6\}$ .

#### Probability

Given 1 trial, in which any root event is expected to occur equally, where  $N$  is the number of all possible events and  $a$  is the number of events  $A$  occurring, then we say  $\frac{a}{N}$  is the **probability** of the event  $A$ , which we express as  $P(A)$ .

$$P(A) = \frac{\text{(Number of cases of event } A \text{ occurring)}}{\text{(Number of all cases that could occur)}} = \frac{a}{N}$$

When we consider multiple combinations, such as of coins or dice, to calculate their probability, we need to distinguish them to find the number of cases.

## EXERCISE



- 1 Given 1 die is rolled, find the probability of an outcome of a 3 or less.

There are a total of 6 possible outcomes.

Furthermore, there are 3 ways, 1, 2, and 3, to roll a 3 or less.

Therefore, we find a probability of  $\frac{3}{6} = \frac{1}{2}$

$\frac{1}{2}$

The content of this page is part of  
"14. Probability".

Students can simulate experiments, such as rolling dice or tossing coins, by using [Math Box]. This allows them to express their opinions through a comparison of experimental probability and theoretical probability.

In **Math Box** on the scientific calculator, you can use the **Dice Roll** or **Coin Toss** simulation to check probability (statistical probability).

Simulate rolling 1 die 250 times.

Press  $\odot$ , select [Math Box], press  $\text{OK}$ , select [Dice Roll], press  $\text{OK}$

Select [Dice], press  $\text{OK}$ , select [1 Die], press  $\text{OK}$



Select [Attempts], press  $\text{OK}$ , after inputting 250 (number of attempts), select [Confirm], press  $\text{OK}$

Select [Same Result], press  $\text{OK}$ , select [Off], press  $\text{OK}$

Select [Execute], press  $\text{EXE}$ , select [Relative Freq], press  $\text{OK}$



Read the values in the table for when the sum of the die was 1, 2, and 3, then

add them to get  $0.18+0.168+0.148=0.496$ , and you can confirm that the

probability is approximately  $0.5 = \frac{1}{2}$ .

(When **Same Result** is Off, the results of each trial are different.)

Sum	Freq	Rel Fr	Attempts
1	45	0.18	250
2	42	0.168	
3	37	0.148	
4	43	0.172	0.148

2 Given 2 coins being tossed at the same time, find the probability of 1 being heads and 1 being tails.

Consider the combinations of heads and tails of the 2 coins.

There are a total of 4 combinations: (Heads, Heads), (Heads, Tails), (Tails, Heads), and (Tails, Tails).

Furthermore, there are 2 ways in which 1 is heads and 1 is tails, as shown above.

Therefore, we find a probability of  $\frac{2}{4} = \frac{1}{2}$

$\frac{1}{2}$

**check**

On the scientific calculator, simulate tossing 2 coins 250 times.

Press  $\odot$ , select [Math Box], press  $\text{OK}$ , select [Coin Toss], press  $\text{OK}$

Select [Coins], press  $\text{OK}$ , select [2 Coins], press  $\text{OK}$

Select [Attempts], press  $\text{OK}$ , after inputting 250, select [Confirm], press  $\text{OK}$

Select [Same Result], press  $\text{OK}$ , select [Off], press  $\text{OK}$

Select [Execute], press  $\text{EXE}$ , select [Relative Freq], press  $\text{OK}$

Side	Freq	Rel Fr	Attempts
•x0	61	0.244	250
•x1	128	0.512	
•x2	61	0.244	0.512

From the table, the value of heads coming up 1 time [ $\bullet \times 1$ ] is about 0.51, and you can confirm that the probability is

approximately  $\frac{1}{2}$ .

# Limit values

## TARGET

To understand limit values and how to find them.

## STUDY GUIDE

### Limit values

In the function  $f(x)$ , when the value  $x$  is infinitely approaching  $a$ , while remaining a different value than  $a$ , then the value of  $f(x)$  is infinitely approaching  $b$ , which is expressed as  $\lim_{x \rightarrow a} f(x) = b$ . This value  $b$  is called the **limit value** of  $f(x)$  when  $x \rightarrow a$ .  $\lim$  is the abbreviated symbol for limit, and is read as "the limit of".

### Limit values

$$\lim_{x \rightarrow a} f(x) = b \text{ or } f(x) \rightarrow b \text{ when } x \rightarrow a$$

**Ex.** In the function  $f(x) = x + 4$ , when the value  $x$  is infinitely approaching 3, the limit value is  $\lim_{x \rightarrow 3} (x + 4) = 7$ .

### EXERCISE



Find the limit values of the following.

(1)  $\lim_{x \rightarrow 5} (x^2 - 3x + 2)$

When the value of  $x$  infinitely approaches 5,  $x^2 - 3x + 2$  approaches  $5^2 - 3 \times 5 + 2 = 12$ .

$$\lim_{x \rightarrow 5} (x^2 - 3x + 2) = 12$$




### check

Use Table (to create a function table) to see how limit values are approached.

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$ .

Press  $\text{f(x)}$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$ , after inputting  $f(x) = x^2 - 3x + 2$ , press  $\text{EXE}$ .

Press  $\text{f(x)}$ , select [Table Range], press  $\text{OK}$ , after inputting [Start:4.9, End:5, Step:0.01], select [Execute], press  $\text{EXE}$ .



	$x$	$f(x)$	$g(x)$
1	4.9	11.378	ERROR
2	4.91	11.378	ERROR
3	4.92	11.446	ERROR
4	4.93	11.514	ERROR

11.31

	$x$	$f(x)$	$g(x)$
5	4.94	11.583	ERROR
6	4.95	11.652	ERROR
7	4.96	11.721	ERROR
8	4.97	11.790	ERROR

11.7909

	$x$	$f(x)$	$g(x)$
8	4.97	11.79	ERROR
9	4.98	11.86	ERROR
10	4.99	11.93	ERROR
11	5	12	ERROR

12

By using [Table], the student can check how the values of the function approaches the limit value.

From the table, we can confirm that the value of  $f(x)$  approaches 12 as the value of  $x$  approaches 5 from 4.9.

In this case, the value approaches 5 while remaining smaller than 5, but we get similar results even though it approaches 5 from the other side.

The content of this page is part of "15. Differential Calculus and Integral Calculus".

$$(2) \lim_{x \rightarrow 2} \frac{3x+10}{x+2}$$

When the value of  $x$  infinitely approaches 2,  $\frac{3x+10}{x+2}$  approaches  $\frac{3 \times 2 + 10}{2+2} = \frac{16}{4} = 4$ .

$$\lim_{x \rightarrow 2} \frac{3x+10}{x+2} = 4$$

4

check

Use Table (to create a function table) to see how limit values are approached.

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

Press  $\ominus$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$ ,

after inputting  $f(x) = \frac{3x+10}{x+2}$ , press  $\text{EXE}$

Press  $\ominus$ , select [Table Range], press  $\text{OK}$ , after inputting

[Start:1.9, End:2, Step:0.01], select [Execute], press  $\text{EXE}$

$x$	$f(x)$	$g(x)$
1	1.9	4.0256
2	1.91	4.023
3	1.92	4.0204
4	1.93	4.0178

$x$	$f(x)$	$g(x)$
5	1.94	4.0152
6	1.95	4.0126
7	1.96	4.0101
8	1.97	4.0075

$$f(x) = \frac{3x+10}{x+2}$$

Table Range  
Start:1.9  
End :2  
Step :0.01

$x$	$f(x)$	$g(x)$
8	1.97	4.0075
9	1.98	4.005
10	1.99	4.0025
11	2	4

From the table, we can confirm that the value of  $f(x)$  approaches 4 as the value of  $x$  approaches 2 from 1.9.

$$(3) \lim_{h \rightarrow 0} \frac{h^2 + 3h}{h}$$

$$\lim_{h \rightarrow 0} \frac{h^2 + 3h}{h} = \lim_{h \rightarrow 0} \frac{h(h+3)}{h} = \lim_{h \rightarrow 0} (h+3)$$

When the value of  $h$  infinitely approaches 0,  $h+3$  approaches  $0+3=3$ .

$$\lim_{h \rightarrow 0} \frac{h^2 + 3h}{h} = 3$$

3

check

Use Table (to create a function table) to see how limit values are approached.

Press  $\odot$ , select [Table], press  $\text{OK}$ , then clear the previous data by pressing  $\text{C}$

Press  $\ominus$ , select [Define  $f(x)/g(x)$ ], press  $\text{OK}$ , select [Define  $f(x)$ ], press  $\text{OK}$ ,

after inputting  $f(x) = \frac{x^2 + 3x}{x}$ , press  $\text{EXE}$

Press  $\ominus$ , select [Table Range], press  $\text{OK}$ , after inputting

[Start:0.1, End:0, Step:-0.01], select [Execute], press  $\text{EXE}$

$x$	$f(x)$	$g(x)$
1	0.1	3.1
2	0.09	3.09
3	0.08	3.08
4	0.07	3.07

$x$	$f(x)$	$g(x)$
5	0.06	3.06
6	0.05	3.05
7	0.04	3.04
8	0.03	3.03

$$f(x) = \frac{x^2 + 3x}{x}$$

Table Range  
Start:0.1  
End :0  
Step :-0.01

$x$	$f(x)$	$g(x)$
8	0.03	3.03
9	0.02	3.02
10	0.01	3.01
11	0	ERROR

From the table, we can confirm that the value of  $f(x)$  approaches 3 as the value of  $x$  approaches 0 from 0.1. In the table,  $x=0$  appears as ERROR because there is no value.

## PRACTICE



- 1 Let  $m$  be the slope of the tangent  $l$  for the point  $P(-1, 9)$  on the graph of the function  $y = -x^2 - 6x + 4$ . Now, solve the following problems.

- (1) Find the value of  $m$ .

**Given  $f(x) = -x^2 - 6x + 4$ , we can get  $m = f'(-1)$ .**

**By differentiating  $f(x)$ , we get  $f'(x) = (-x^2 - 6x + 4)' = -2x - 6$ .**

**Therefore, we get  $m = f'(-1) = -2 \cdot (-1) - 6 = -4$ .**

Scientific calculators can also be used for calculations to find differential coefficients.

⊞ OK OK − (X) x<sup>2</sup> − 6 (X) + 4 > − 1 EXE

$$\left. \frac{d}{dx}(-x^2 - 6x + 4) \right|_{x=-1} = -4$$

−4

- (2) Find the equation of tangent  $l$ .

**The tangent  $l$  is a straight line passing through the point  $P(-1, 9)$  with a slope of  $m = -4$ .**

**Therefore, from  $y - 9 = -4\{x - (-1)\}$ , we get  $y = -4x + 5$ .**

$$y = -4x + 5$$

check

**To better understand, use Table and the QR code to calculate and draw the relation of the tangent and graph of the function.**

**Press ⊞, select [Table], press OK, then clear the previous data by pressing ⊞**

**Press ⊞, select [Define f(x)/g(x)], press OK, select [Define f(x)], press OK, after inputting  $f(x) = -x^2 - 6x + 4$ , press EXE**

**In the same way, input  $g(x) = -4x + 5$ .**

**Press ⊞, select [Table Range], press OK, after inputting [Start:−5, End:2, Step:1], select [Execute], press EXE**

**Press ↑ (X), scan the QR code to display a graph.**

$$f(x) = -x^2 - 6x + 4$$

$$g(x) = -4x + 5$$


Table Range  
Start:−5  
End :2  
Step :1

x	f(x)	g(x)
−5	9	25
−4	12	21
−3	13	17
−2	12	13
−1	9	9
0	4	5
1	−3	1
2	−12	−3

x	f(x)	g(x)
−5	9	25
−4	12	21
−3	13	17
−2	12	13
−1	9	9
0	4	5
1	−3	1
2	−12	−3

**From the table and graph, we can see that the maximum value for  $f(x)$  is  $x = -3$ .**

**$f(x)$  and  $g(x)$  share a point  $(-1, 9)$ .**



The content of this page is part of "15. Differential Calculus and Integral Calculus".

**CASIO Teaching Materials**  
**Basic Mathematics Exercise with fx-991CW**  
**DIGEST EDITION**

Publisher: CASIO COMPUTER CO., Ltd.

Date of Publication: 2025/5/31 (1st edition)

<https://edu.casio.com>

Copyright © 2025 CASIO COMPUTER CO., LTD.

(1) These teaching materials may be freely reproduced and distributed by teachers for educational purposes.

\* However, publication on the Web or simultaneous transmission to an unspecified number of people is prohibited.

(2) Any reproduction, distribution, editing, or use for purposes other than those listed above requires the permission of CASIO COMPUTER CO., LTD.

(3) QR Code is a registered trademark of DENSO WAVE INCORPORATED in Japan and in other countries.



**Boost  
your  
Curiosity**